

CORRIGENDUM: A MIXED BOOLEAN AND DEPOSIT MODEL FOR THE MODELING OF METAL PIGMENTS IN PAINT LAYERS (2015 IMAGE ANAL STEREOLOG 34:125–34)

ENGUERRAND COUKA, FRANÇOIS WILLOT AND DOMINIQUE JEULIN

Center for Mathematical Morphology, MINES ParisTech, PSL Research University, 35 rue Saint-Honoré, 77300 Fontainebleau, France

e-mail: enguerrand.couka@mines-paristech.fr, francois.willot@mines-paristech.fr,
 dominique.jeulin@mines-paristech.fr

(Accepted October 27, 2015)

The expressions for the distribution function and the cumulative distribution of chord lengths given in Couka *et al.* (2015) (Eqs. 3 and 4) are incorrect. The latter must be replaced by:

$$P_1(\ell) = P\{L < \ell\} = 1 - \sqrt{1 - \left(\frac{\ell}{D}\right)^2}, \quad (1)$$

$$P_2(\ell) = 1 - \frac{\ell}{D} \left[\sqrt{\left(\frac{D}{\ell}\right)^2 - 1} - \cos^{-1}\left(\frac{\ell}{D}\right) \right], \quad (2)$$

respectively. The above expressions are numerically very similar to that given in Couka *et al.* (2015) and to the measurements carried out on the SEM images (see updated Figs. 1a and b, which replace Figs. 7 and 18, resp).




Fig. 1. (a): Cumulative distribution of the lengths of the particles along their main direction, on a 2D section: measurements (red), theoretical distribution for cylinders with constant (green) or uniform diameter (blue). (b): Cumulative distribution function of the length of particles along the main directions as observed on SEM images (blue) and stack model (red). Theoretical formula seen in Eq. (2) shown in green.

REFERENCES

Couka E, Willot F, Jeulin D (2015). A mixed Boolean and deposit model for the modeling of metal pigments in paint layers. *Image Anal Stereol* 34(2):125–34.